arXiv:1201.5494v2 [math.CA] 7 Feb 2012

# Discontinuous Sturm-Liouville type problems

Erdoğan Şen

Department of Mathematics, Faculty of Science and Letters, Namık Kemal University, 59030 Tekirdağ, Turkey.

e-mail: esen@nku.edu.tr

**MSC (2010):** 34L20, 35R10.

**Keywords :** Sturm-Liouville problems; Transmission conditions; Asymptotics of eigenvalues and eigenfunctions.

ABSTRACT.

In this work, we study Sturm-Liouville problems with eigenparameter dependent boundary condition and transmission conditions at three interior points. We obtain asymptotic formulae for the eigenvalues and eigenfunctions.

## 1 Introduction

It is well-known that many topics in mathematical physics require the investigation of eigenvalues and eigenfunctions of Sturm-Liouville type boundary value problems. In recent years, more and more researches are interested in the discontinuous Sturm-Liouville problem for its applications in physics (see $[1-4]$). Various physics applications of this kind problem are found in many literatures, including some boundary value problem with transmission conditions that arise in the theory of heat and mass transfer (see $[5-9]$). The literature on such results is voluminous and we refer to $[7, 8, 10-20]$.

Discontinuous Sturm-Liouville problems with eigen-dependent boundary conditions and with two supplementary transmission conditions at the point(s) of discontinuity have been investigated in [10, 12, 21, 22]. Note that discontinuous Sturm-Liouville problems with eigen-dependent boundary conditions and with four supplementary transmission conditions at the points of discontinuity have been investigated in [3].

In this study, we shall consider discontinuous eigenvalue problem which consists of Sturm-Liouville equation

$$Lu := -u'' + q(x)u = \lambda u \tag{1.1}$$

on $[a, h_1) \cup (h_1, h_2) \cup (h_2, h_3) \cup (h_3, b]$, with boundary condition at $x = a$

$$L_1 u := \alpha_1 u\left(a\right) + \alpha_2 u^{'}\left(a\right) = 0, \tag{1.2}$$

with the six transmission conditions at the points of discontinuity $x = h_1, x = h_2$ and $x = h_3$,

$$L_2 u := \gamma_1 u\left(h_1 - 0\right) - \delta_1 u\left(h_1 + 0\right) = 0, \tag{1.3}$$

$$L_3 u := \gamma_1^{'} u^{'}\left(h_1 - 0\right) - \delta_1^{'} u^{'}\left(h_1 + 0\right) = 0, \tag{1.4}$$

$$L_4 u := \gamma_2 u\left(h_2 - 0\right) - \delta_2 u\left(h_2 + 0\right) = 0, \tag{1.5}$$

$$L_5 u := \gamma_2^{'} u^{'}\left(h_2 - 0\right) - \delta_2^{'} u^{'}\left(h_2 + 0\right) = 0, \tag{1.6}$$

$$L_6 u := \gamma_3 u\left(h_3 - 0\right) - \delta_3 u\left(h_3 + 0\right) = 0, \tag{1.7}$$

$$L_7 u := \gamma_3^{'} u^{'}\left(h_3 - 0\right) - \delta_3^{'} u^{'}\left(h_3 + 0\right) = 0 \tag{1.8}$$

and the eigen-dependent boundary condition at $x = b$

$$L_8 u := \lambda\left[\beta_1^{'} u\left(b\right) - \beta_2^{'} u^{'}\left(b\right)\right] + \left[\beta_1 u\left(b\right) - \beta_2 u^{'}\left(b\right)\right] = 0 \tag{1.9}$$

where $q(x)$ is a given real-valued function continuous in $\left[a, h_1\right), \left(h_1, h_2\right), \left(h_2, h_3\right)$ and $\left(h_3, b\right]$ and has finite limits $q(h_1\pm) = \lim_{x\to h_1} \pm q(x)$, $q(h_2\pm) = \lim_{x\to h_2} \pm q(x)$, $q(h_3\pm) = \lim_{x\to h_3} \pm q(x)$; $\lambda$ is a complex eigenvalue parameter; the coefficients of the boundary and transmission conditions are real numbers. We assume $|a_1| + |a_2| \neq 0$, $|\gamma_i| + |\delta_i| \neq 0$, $|\gamma_i| + |\delta_i| \neq 0$ $(i = 1, 2, 3)$ and $\rho = \begin{vmatrix} \beta_1^{'} & \beta_2' \\ \beta_3^{'} & \beta_4' \end{vmatrix} > 0$. Contrary to previous works, the eigenfunctions of this problem may have discontinuities.

## 2 Preliminaries

For the sake of convenience let us introduce the following notations:

$$\Omega_1 := \left[a, h_1\right], \ \Omega_2 := \left[h_1, h_2\right], \ \Omega_3 := \left[h_2, h_3\right], \ \Omega_4 := \left[h_3, b\right]$$

$$u_{(1)}\left(x\right) := \begin{cases} u(x) & x \in \left[a, h_1\right) \\ \lim_{x\to h_1 -} u(x) & x = h_1, \end{cases} \quad u_{(2)}\left(x\right) := \begin{cases} u(x) & x \in \left(h_1, h_2\right) \\ \lim_{x\to h_1 +} u(x) & x = h_1, \end{cases}$$

$$u_{(3)}\left(x\right) := \begin{cases} u(x) & x \in \left(h_1, h_2\right) \\ \lim_{x\to h_2 -} u(x) & x = h_2, \end{cases} \quad u_{(4)}\left(x\right) := \begin{cases} u(x) & x \in \left(h_2, h_3\right) \\ \lim_{x\to h_2 +} u(x) & x = h_2, \end{cases}$$

$$u_{(5)}\left(x\right) := \begin{cases} u(x) & x \in \left(h_2, h_3\right) \\ \lim_{x\to h_3 -} u(x) & x = h_3, \end{cases} \quad u_{(6)}\left(x\right) := \begin{cases} u(x) & x \in \left(h_3, b\right] \\ \lim_{x\to h_3 +} u(x) & x = h_3, \end{cases}$$

$$\left(u\right)_\beta := \lim_{x\to b}\left(\beta_1 u\left(x\right) - \beta_2 u^{'}\left(x\right)\right), \quad \left(u\right)_\beta^{'} := \lim_{x\to b}\left(\beta_1^{'} u\left(x\right) - \beta_2^{'} u^{'}\left(x\right)\right),$$

$$\tilde{u}\left(x\right) := \begin{cases} u(x) & x \in \left[a, b\right) \\ \left(u\right)_\beta^{'} & x = b. \end{cases}$$

Note that everywhere below, we shall assume that $\gamma_i \gamma_i^{'} \delta_i \delta_i^{'} > 0$ $(i = 1, 2, 3)$ and for the Lebesque measurable subsets $M \subset \left[a, h_1\right) \cup \left(h_1, h_2\right) \cup \left(h_2, h_3\right) \cup \left(h_3, b\right]$ with Lebesque measure $\mu_L\left(M\right)$, we shall define a new positive measure $\mu_\rho\left(M\right)$

by

$$\mu_\rho(M) := \frac{\gamma_1\gamma_1^{'}}{\delta_1\delta_1^{'}}\mu_L(M\cap[a,h_1)) + \mu_L(M\cap(h_1,h_2)) + \frac{\delta_2\delta_2^{'}}{\gamma_2\gamma_2^{'}}\mu_L(M\cap(h_2,h_3))$$
$$\frac{\delta_2\delta_2^{'}\delta_3\delta_3^{'}}{\gamma_2\gamma_2^{'}\gamma_3\gamma_3^{'}}\mu_L(M\cap(h_3,b]) + \frac{\delta_2\delta_2^{'}\delta_3\delta_3^{'}}{\gamma_2\gamma_2^{'}\gamma_3\gamma_3^{'}}\frac{b(M)}{\rho}$$

where

$$b(M) := \begin{cases} 0, & \text{if } b\notin M,\\ 1, & \text{if } b\in M.\end{cases}$$

Let $\langle\cdot,\cdot\rangle_{H_\rho}$ denote the scalar product in the Hilbert space $H_\rho := \left(L^2[a,b];\mu_\rho\right)$. In this space, we define a linear operator $A$ by the domain of definition

$$D(A) := \Big\{u\in H_\rho \,\Big|\, u_{(i)}, u_{(i)}^{'} \text{ are absolutely continuous in } \Omega_i \;\; (i=1,2,3,4),\ L\widetilde{u}\in L^2[a,b]$$
$$\alpha_1\widetilde{u}(a) + \alpha_2\widetilde{u}^{'}(a) = 0,\ \gamma_1\widetilde{u}(h_1-0) = \delta_1\widetilde{u}(h_1+0),\ \gamma_1^{'}\widetilde{u}^{'}(h_1-0) = \delta_1^{'}\widetilde{u}^{'}(h_1+0)$$
$$\gamma_2\widetilde{u}(h_2-0) = \delta_2\widetilde{u}(h_2+0),\ \gamma_2^{'}\widetilde{u}^{'}(h_2-0) = \delta_2^{'}\widetilde{u}^{'}(h_2+0),\ \gamma_3\widetilde{u}(h_3-0) = \delta_3\widetilde{u}(h_3+0)$$
$$\gamma_3^{'}\widetilde{u}^{'}(h_3-0) = \delta_3^{'}\widetilde{u}^{'}(h_3+0)\Big\}$$

and

$$(A\widetilde{u})(x) = \begin{cases} (L\widetilde{u})(x) & \text{for } x\in[a,h_1)\cup(h_1,h_2)\cup(h_2,h_3)\cup(h_3,b] \\ -(\widetilde{u})_\beta & \text{, for } x=b\end{cases}$$

Consequently, the considered problem (1.1)-(1.9) can be rewritten in operator form as

$$A\widetilde{u} = \lambda\widetilde{u}$$

i.e., the problem (1.1)-(1.9) can be considered as the eigenvalue problem for the operator $A$.

**Theorem 2.1.** *The operator $A$ is symmetric.*

**Proof.** Let $f\ , g\in D(A)$. By the means of two partial integrations, we get

$$\langle Af,g\rangle_{H_\rho} - \langle f,Ag\rangle_{H_\rho} = \frac{\gamma_1\gamma_1^{'}}{\delta_1\delta_1^{'}}\left(W(f,\overline{g};h_1-0) - W(f,\overline{g};a)\right) + \left(W(f,\overline{g};h_2-0) - W(f,\overline{g};h_1+0)\right)$$
$$\frac{\delta_2\delta_2^{'}}{\gamma_2\gamma_2^{'}}\left(W(f,\overline{g};h_3-0) - W(f,\overline{g};h_2+0)\right) + \frac{\delta_2\delta_2^{'}\delta_3\delta_3^{'}}{\gamma_2\gamma_2^{'}\gamma_3\gamma_3^{'}}$$
$$\times\left(W(f,\overline{g};b) - W(f,\overline{g};h_3+0)\right) + \frac{\delta_2\delta_2^{'}\delta_3\delta_3^{'}}{\rho\gamma_2\gamma_2^{'}\gamma_3\gamma_3^{'}}\left((f\ )_\beta^{'}(\overline{g})_\beta - (f\ )_\beta(\overline{g})_\beta^{'}\right) \tag{2.1}$$

where, as usual,

$$W(f,g;x) = f(x)\,g^{'}(x) - f^{'}(x)\,g(x) \tag{2.2}$$

denotes the Wronskians of the functions $f$ and $g$. Since $f$ and $\overline{g}$ satisfy the boundary condition (1.2), it follows that

$$W(f,g;a) = 0. \tag{2.3}$$

From the transmission conditions (1.3)-(1.8) we get

$$\gamma_i\gamma_i^{'}W\left(f,g;h_i-0\right)=\delta_i\delta_i^{'}W\left(f,g;h_i+0\right) \tag{2.4}$$

Furthermore, it is easy to verify that

$$\left(f\ \right)_\beta\left(\overline{g}\right)_\beta^{'}-\left(f\ \right)_\beta^{'}\left(\overline{g}\right)_\beta=\rho W\left(f,g;b\right). \tag{2.5}$$

Finally, substituting (2.2)-(2.5) in (2.1) yields the required equality

$$\left\langle Af,g\right\rangle_{H_\rho}=\left\langle f,Ag\right\rangle_{H_\rho}\quad\left(f,g\in H_\rho\right). \tag{2.6}$$

■

**Corollary 2.1.** *All eigenvalues of the problem (1.1)-(1.9) are real.*

We can now assume that all eigenfunctions are real-valued.

**Corollary 2.2.** *If $\lambda_1$ and $\lambda_2$ are two different eigenvalues of the problem (1.1)-(1.9), then corresponding eigenfunctions $u_1$ and $u_2$ of this problem satisfy the following equality*:

$$\frac{\gamma_1\gamma_1^{'}}{\delta_1\delta_1^{'}}\int_a^{h_1}u_1\left(x\right)u_2\left(x\right)dx+\int_{h_1}^{h_2}u_1\left(x\right)u_2\left(x\right)dx+\frac{\delta_2\delta_2^{'}}{\gamma_2\gamma_2^{'}}\int_{h_2}^{h_3}u_1\left(x\right)u_2\left(x\right)dx$$

$$\frac{\delta_2\delta_2^{'}\delta_3\delta_3^{'}}{\gamma_2\gamma_2^{'}\gamma_3\gamma_3^{'}}\int_{h_3}^{b}u_1\left(x\right)u_2\left(x\right)dx+\frac{\delta_2\delta_2^{'}\delta_3\delta_3^{'}}{\gamma_2\gamma_2^{'}\gamma_3\gamma_3^{'}}\left(u_1\right)_\beta^{'}\left(u_2\right)_\beta^{'}=0. \tag{2.7}$$

*In fact this formula means the orthogonality of eigenfunctions $u_1$ and $u_2$ in the Hilbert space $H_\rho$.*

We need the following lemma, which can be proved similarly to [10, Theorem 2].

**Lemma 2.1.** *Let the real-valued function $q\left(x\right)$ be continuous in $[a,b]$ where $f\left(\lambda\right),g\left(\lambda\right)$ are given entire functions. Then for any $\lambda\in\mathbb{C}$ the equation*

$$-u^{''}+q\left(x\right)u=\lambda u,\quad x\in[a,b]$$

*has a unique solution $u=u\left(x,\lambda\right)$ such that*

$$u\left(a\right)=f\left(\lambda\right),u^{'}\left(a\right)=g\left(\lambda\right)\ \left(\text{or }u\left(b\right)=f\left(\lambda\right),u^{'}\left(b\right)=g\left(\lambda\right)\right)$$

*and for each $x\in[a,b]$, $u\left(x,\lambda\right)$ is an entire function of $\lambda$.*

We shall define two solutions

$$\phi_\lambda\left(x\right)=\begin{cases}\phi_{1\lambda}\left(x\right),\ x\in[a,h_1),\\ \phi_{2\lambda}\left(x\right),\ x\in(h_1,h_2),\\ \phi_{3\lambda}\left(x\right),\ x\in(h_2,h_3),\\ \phi_{4\lambda}\left(x\right),\ x\in(h_3,b],\end{cases}\quad\text{and }\chi_\lambda\left(x\right)=\begin{cases}\chi_{1\lambda}\left(x\right),\ x\in[a,h_1),\\ \chi_{2\lambda}\left(x\right),\ x\in(h_1,h_2),\\ \chi_{3\lambda}\left(x\right),\ x\in(h_2,h_3),\\ \chi_{4\lambda}\left(x\right),\ x\in(h_3,b],\end{cases}$$

of the Eq. (1.1) as follows: Let $\phi_{1\lambda}\left(x\right)=\phi_1\left(x,\lambda\right)$ be the solution of Eq. (1.1) on $[a,h_1]$, which satisfies the initial conditions

$$\begin{pmatrix}u\left(a\right)\\ u^{'}\left(a\right)\end{pmatrix}=\begin{pmatrix}\alpha_2\\ -\alpha_1\end{pmatrix}. \tag{2.8}$$

By virtue of Lemma 2.1, after defining this solution, we may define the solution $\phi_2(x,\lambda)$ of Eq. (1.1) on $[h_1,h_2]$ by means of the solution $\phi_1(x,\lambda)$ by the nonstandart initial conditions

$$\left(\begin{array}{c} u(h_1+0) \\ u^{'}(h_1+0) \end{array}\right) = \left(\begin{array}{c} \frac{\gamma_1}{\delta_1}\phi_{1\lambda}(h_1-0,\lambda) \\ \frac{\gamma_1^{'}}{\delta_1^{'}}\phi_{1\lambda}^{'}(h_1-0,\lambda) \end{array}\right). \tag{2.9}$$

After defining this solution, we may define the solution $\phi_3(x,\lambda)$ of Eq. (1.1) on $[h_2,h_3]$ by means of the solution $\phi_2(x,\lambda)$ by the nonstandart initial conditions

$$\left(\begin{array}{c} u(h_2+0) \\ u^{'}(h_2+0) \end{array}\right) = \left(\begin{array}{c} \frac{\gamma_2}{\delta_2}\phi_{2\lambda}(h_2-0,\lambda) \\ \frac{\gamma_2^{'}}{\delta_2^{'}}\phi_{2\lambda}^{'}(h_2-0,\lambda) \end{array}\right). \tag{2.10}$$

After defining this solution, we may define the solution $\phi_4(x,\lambda)$ of Eq. (1.1) on $[h_3,b]$ by means of the solution $\phi_3(x,\lambda)$ by the nonstandart initial conditions

$$\left(\begin{array}{c} u(h_3+0) \\ u^{'}(h_3+0) \end{array}\right) = \left(\begin{array}{c} \frac{\gamma_3}{\delta_3}\phi_{3\lambda}(h_3-0,\lambda) \\ \frac{\gamma_3^{'}}{\delta_3^{'}}\phi_{3\lambda}^{'}(h_3-0,\lambda) \end{array}\right). \tag{2.11}$$

Therefore, $\phi(x,\lambda)$ satisfies the Eq. (1.1) on $[a,h_1)\cup(h_1,h_2)\cup(h_2,h_3)\cup(h_3,b]$, the boundary condition (1.2), and the transmission conditions (1.3)-(1.8).

Similarly, first we define the solution $\chi_{4\lambda}(x) := \chi_4(x,\lambda)$ on $[h_3,b]$ by the initial conditions

$$\left(\begin{array}{c} u(b) \\ u^{'}(b) \end{array}\right) = \left(\begin{array}{c} \beta_2^{'}\lambda+\beta_2 \\ \beta_1^{'}\lambda+\beta_1 \end{array}\right). \tag{2.12}$$

Again, after defining this solution, we may define the solution $\chi_{3\lambda}(x) := \chi_3(x,\lambda)$ of the Eq. (1.1) on $[h_2,h_3]$ by the initial conditions

$$\left(\begin{array}{c} u(h_3-0) \\ u^{'}(h_3-0) \end{array}\right) = \left(\begin{array}{c} \frac{\delta_3}{\gamma_3}\chi_{4\lambda}(h_3+0,\lambda) \\ \frac{\delta_3^{'}}{\gamma_3^{'}}\chi_{4\lambda}^{'}(h_3+0,\lambda) \end{array}\right). \tag{2.13}$$

After defining this solution, we may define the solution $\chi_{2\lambda}(x) := \chi_2(x,\lambda)$ of the Eq. (1.1) on $[h_1,h_2]$ by the initial conditions

$$\left(\begin{array}{c} u(h_2-0) \\ u^{'}(h_2-0) \end{array}\right) = \left(\begin{array}{c} \frac{\delta_2}{\gamma_2}\chi_{3\lambda}(h_2+0,\lambda) \\ \frac{\delta_2^{'}}{\gamma_2^{'}}\chi_{3\lambda}^{'}(h_2+0,\lambda) \end{array}\right). \tag{2.14}$$

After defining this solution, we may define the solution $\chi_{1\lambda}(x) := \chi_1(x,\lambda)$ of the Eq. (1.1) on $[a,h_1]$ by the initial conditions

$$\left(\begin{array}{c} u(h_1-0) \\ u^{'}(h_1-0) \end{array}\right) = \left(\begin{array}{c} \frac{\delta_1}{\gamma_1}\chi_{2\lambda}(h_1+0,\lambda) \\ \frac{\delta_1^{'}}{\gamma_1^{'}}\chi_{2\lambda}^{'}(h_1+0,\lambda) \end{array}\right). \tag{2.15}$$

Therefore, $\chi(x,\lambda)$ satisfies the Eq. (1.1) on $[a,h_1)\cup(h_1,h_2)\cup(h_2,h_3)\cup(h_3,b]$, the boundary condition (1.9), and the transmission conditions (1.3)-(1.8).

Further it follows from (1.1) that the Wronskians

$$\omega_i := W_\lambda(\phi_i,\chi_i;x) := \phi_i(x,\lambda)\chi_i^{'}(x,\lambda) - \phi_i^{'}(x,\lambda)\chi_i(x,\lambda), \quad x\in\Omega_i \ \ (i=1,2,3,4)$$

are independent of $x\in\Omega_i$. Moreover these functions are entire functions of $\lambda$.

**Lemma 2.2.** *For each* $\lambda\in\mathbb{C}$,

$$\gamma_1\gamma_1^{'}\gamma_2\gamma_2^{'}\gamma_3\gamma_3^{'}\omega_1(\lambda) = \delta_1\delta_1^{'}\gamma_2\gamma_2^{'}\gamma_3\gamma_3^{'}\omega_2(\lambda) = \delta_1\delta_1^{'}\delta_2\delta_2^{'}\gamma_3\gamma_3^{'}\omega_3(\lambda) = \delta_1\delta_1^{'}\delta_2\delta_2^{'}\delta_3\delta_3^{'}\omega_4(\lambda).$$

**Proof.** Using (2.9), (2.10), (2.11), (2.13), (2.14) and (2.15), we get

$$\gamma_1\gamma_1^{'}\gamma_2\gamma_2^{'}\gamma_3\gamma_3^{'}W(\phi_1,\chi_1;h_1-0) = \gamma_2\gamma_2^{'}\gamma_3\gamma_3^{'}\delta_1\delta_1^{'}W(\phi_2,\chi_2;h_1+0) =$$
$$\delta_2\delta_2^{'}\gamma_3\gamma_3^{'}\delta_1\delta_1^{'}W(\phi_3,\chi_3;h_2+0) = \delta_1\delta_1^{'}\delta_2\delta_2^{'}\delta_3\delta_{31}^{''}W(\phi_4,\chi_4;h_3+0)$$

which completes the proof. ■

Now we may introduce the characteristic function

$$\begin{aligned}\omega(\lambda) &:= \gamma_1\gamma_1^{'}\gamma_2\gamma_2^{'}\gamma_3\gamma_3^{'}\omega_1(\lambda) = \delta_1\delta_1^{'}\gamma_2\gamma_2^{'}\gamma_3\gamma_3^{'}\omega_2(\lambda)\\ &= \delta_1\delta_1^{'}\delta_2\delta_2^{'}\gamma_3\gamma_3^{'}\omega_3(\lambda) = \delta_1\delta_1^{'}\delta_2\delta_2^{'}\delta_3\delta_3^{'}\omega_4(\lambda).\end{aligned}$$

**Theorem 2.2.** *The eigenvalues of the problem (1.1)-(1.9) are the zeros of the function* $\omega(\lambda)$.

**Proof.** Let $\omega(\lambda_0)=0$. Then $W_{\lambda_0}(\phi_1,\chi_1;x)=0$ and therefore the functions $\phi_{1\lambda_0}(x)$ and $\chi_{1\lambda_0}(x)$ are linearly dependent, i.e.

$$\chi_{1\lambda_0}(x) = k_1\phi_{1\lambda_0}(x), \quad x\in[a,h_1]$$

for some $k_1\neq 0$. From this, it follows that $\chi(x,\lambda_0)$ satisfies also the first boundary condition (1.2), so $\chi(x,\lambda_0)$ is an eigenfunction for the eigenvalue $\lambda_0$.

Now we let $u_0(x)$ be any eigenfunction corresponding to eigenvalue $\lambda_0$, but $\omega(\lambda_0)\neq 0$. Then the pair of the functions $(\phi_1,\chi_1)$, $(\phi_2,\chi_2)$, $(\phi_3,\chi_3)$ and $(\phi_4,\chi_4)$ would be linearly independent on $[a,h_1)$, $(h_1,h_2)$, $(h_2,h_3)$ and $(h_3,b]$ respectively. Therefore $u_0(x)$ may be represented as

$$u_0(x) = \begin{cases} c_1\phi_1(x,\lambda_0) + c_2\chi_1(x,\lambda_0), & x\in[a,h_1), \\ c_3\phi_2(x,\lambda_0) + c_4\chi_2(x,\lambda_0), & x\in(h_1,h_2), \\ c_5\phi_3(x,\lambda_0) + c_6\chi_3(x,\lambda_0), & x\in(h_2,h_3), \\ c_7\phi_4(x,\lambda_0) + c_8\chi_4(x,\lambda_0), & x\in(h_3,b]. \end{cases}$$

where at least one of the constants $c_1$, $c_2$, $c_3$, $c_4$, $c_5$, $c_6$, $c_7$ and $c_8$ is not zero. Considering the equations

$$\tau_\upsilon(u_0(x)) = 0, \quad \upsilon=\overline{1,8} \tag{2.16}$$

as a system of linear equations of the variables $c_1$, $c_2$, $c_3$, $c_4$, $c_5$, $c_6$, $c_7$, $c_8$ and taking (2.9), (2.10), (2.11), (2.13), (2.14) and (2.15) into account, it follows that the determinant of this system is

$$\begin{vmatrix} S_1 & S_2 \\ S_3 & S_4 \end{vmatrix} = -\delta_1\delta_1^{'}\delta_2\delta_2^{'}\delta_3\delta_3^{'}\omega_1(\lambda_0)\,\omega_2(\lambda_0)\,\omega_3(\lambda_0)\,\omega^4(\lambda_0)\neq 0,$$

where

$$S_1 = \begin{bmatrix} 0 & \omega_1(\lambda_0) & 0 & 0 \\ \gamma_1\phi_{1\lambda_0}(h_1-0) & \gamma_1\chi_{1\lambda_0}(h_1-0) & -\delta_1\phi_{2\lambda_0}(h_1+0) & -\delta_1\chi_{2\lambda_0}(h_1+0) \\ \gamma_1^{'}\phi_{1\lambda_0}^{'}(h_1-0) & \gamma_1^{'}\chi_{1\lambda_0}^{'}(h_1-0) & -\delta_1^{'}\phi_{2\lambda_0}^{'}(h_1+0) & -\delta_1^{'}\chi_{2\lambda_0}^{'}(h_1+0) \\ 0 & 0 & \gamma_2\phi_{2\lambda_0}(h_2-0) & \gamma_2\chi_{2\lambda_0}(h_2-0) \end{bmatrix},$$

$$S_2 = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -\delta_2\phi_{3\lambda_0}(h_2+0) & -\delta_2\chi_{3\lambda_0}(h_2+0) & 0 & 0 \end{bmatrix},$$

$$S_3 = \begin{bmatrix} 0 & 0 & \gamma_2^{'}\phi_{2\lambda_0}^{'}(h_2-0) & \gamma_2^{'}\chi_{2\lambda_0}^{'}(h_2-0) \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

and

$$S_4 = \begin{bmatrix} -\delta_2^{'}\phi_{3\lambda_0}^{'}(h_2+0) & \delta_2^{'}\chi_{3\lambda_0}^{'}(h_2+0) & 0 & 0 \\ \gamma_3\phi_{3\lambda_0}(h_3-0) & \gamma_3\chi_{3\lambda_0}(h_3-0) & -\delta_3\phi_{4\lambda_0}(h_3+0) & -\delta_3\chi_{4\lambda_0}(h_3+0) \\ \gamma_3^{'}\phi_{3\lambda_0}^{'}(h_3-0) & \gamma_3^{'}\chi_{3\lambda_0}^{'}(h_3-0) & -\delta_3^{'}\phi_{4\lambda_0}^{'}(h_3+0) & -\delta_3^{'}\chi_{4\lambda_0}^{'}(h_3+0) \\ 0 & 0 & \omega_4(\lambda_0) & 0 \end{bmatrix}.$$

Therefore, the system (2.16) has only the trivial solution $c_i = 0$ $\left(i = \overline{1,8}\right)$. Thus we get a contradiction, which completes the proof. ■

**Lemma 2.3.** *If* $\lambda = \lambda_0$ *is an eigenvalue, then* $\phi(x,\lambda_0)$ *and* $\chi(x,\lambda_0)$ *are linearly dependent.*

**Proof.** Let $\lambda = \lambda_0$ be an eigenvalue. Then by virtue of Theorem 2.2

$$W(\phi_{i\lambda_0}, \chi_{i\lambda_0}; x) = \omega_i(\lambda_0) = 0$$

and hence

$$\chi_{i\lambda_0}(x) = k_i\phi_{i\lambda_0}(x) \qquad (i = 1,2,3,4) \tag{2.17}$$

for some $k_1 \neq 0$, $k_2 \neq 0$, $k_3 \neq 0$, and $k_4 \neq 0$, We must show that $k_1 = k_2 = k_3 = k_4$. Suppose, if possible, that $k_1 \neq k_2$.

Taking into account the definitions of the solutions $\phi_i(x,\lambda)$ and $\chi_i(x,\lambda)$ and the equalities (2.17), we have

$$\begin{aligned} \delta_1(k_1-k_2)\phi_{2\lambda}(h_1+0) &= \delta_1 k_1\phi_{2\lambda}(h_1+0) - \delta_1 k_2\phi_{2\lambda}(h_1+0) \\ &= k_1\gamma_1\phi_{1\lambda}(h_1-0) - \delta_1 k_2\phi_{2\lambda}(h_1+0) \\ &= \gamma_1\chi_{1\lambda}(h_1-0) - \delta_1\chi_{2\lambda}(h_1+0) = 0. \end{aligned}$$

Therefore

$$\phi_{2\lambda_0}(h_1+0) = 0. \tag{2.18}$$

Similarly starting from $\delta_1^{'}(k_1-k_2)\phi_{2\lambda}^{!}(h_1+0)$ and following the same procedure, we can derive that

$$\phi_{2\lambda_0}^{'}(h_1+0) = 0. \tag{2.19}$$

From the fact that $\phi_{2\lambda_0}$ is a solution of the differential equation (1.1) on $[h_1, h_2]$ and satisfies the initial conditions (2.18) and (2.19), it follows that $\phi_{2\lambda}(x) = 0$ identically on $[h_1, h_2]$. Making use of (2.9), (2.10), (2.11), (2.18) and (2.19), we may also derive that

$$\phi_{1\lambda_0}(h_1 - 0) = \phi'_{1\lambda_0}(h_1 - 0) = 0,$$
$$\phi_{3\lambda_0}(h_2 + 0) = \phi'_{3\lambda_0}(h_2 + 0) = 0$$

and

$$\phi_{4\lambda_0}(h_3 + 0) = \phi'_{4\lambda_0}(h_3 + 0) = 0$$

respectively. From this, by the same argument as for $\phi_{2\lambda_0}(x)$, it follows that $\phi_{1\lambda_0}(x) = 0$ identically on $[a, h_1]$, $\phi_{3\lambda_0}(x) = 0$ identically on $[h_2, h_3]$ and $\phi_{4\lambda_0}(x) = 0$ identically on $[h_3, b]$. Hence $\phi(x, \lambda_0) = 0$ identically on $[a, h_1) \cup (h_1, h_2) \cup (h_2, h_3) \cup (h_3, b]$. But this contradicts (2.8), since $|\alpha_1| + |\alpha_2| \neq 0$. ■

**Corollary 2.3.** *If $\lambda = \lambda_0$ is an eigenvalue, then both $\phi(x, \lambda_0)$ and $\chi(x, \lambda_0)$ are eigenfunctions corresponding to this eigenvalue.*

**Lemma 2.4.** *All eigenvalues $\lambda_n$ are simple zeros of $\omega(\lambda)$.*

**Proof.** Using the Lagrange's formula (cf. [23], p. 6-7), it can be shown that

$$(\lambda - \lambda_n)\left(\int_a^{h_1} \phi_\lambda(x)\phi_{\lambda_n}(x)\,dx + \int_{h_1}^{h_2} \phi_\lambda(x)\phi_{\lambda_n}(x)\,dx + \int_{h_2}^{h_3} \phi_\lambda(x)\phi_{\lambda_n}(x)\,dx + \int_{h_3}^{b} \phi_\lambda(x)\phi_{\lambda_n}(x)\,dx\right)$$
$$= W(\phi_\lambda, \phi_{\lambda_n}; b) \qquad (2.20)$$

for any $\lambda$. Recall that

$$\chi_{\lambda_n}(x) = k_n \phi_{\lambda_n}(x), \quad x \in [a, h_1) \cup (h_1, h_2) \cup (h_2, h_3) \cup (h_3, b]$$

for some $k_n \neq 0$, $n = 1, 2, ....$ Using this equality for the right side of (2.20), we have

$$W(\phi_\lambda, \phi_{\lambda_n}; b) = \frac{1}{k_n} W(\phi_\lambda, \chi_{\lambda_n}; b) = \frac{1}{k_n}\left(\lambda_n (\phi_\lambda)'_\beta + (\phi_\lambda)_\beta\right) = \frac{1}{k_n}\left[\omega(\lambda) + (\lambda - \lambda_n)(\phi_\lambda)'_\beta\right]$$
$$= (\lambda - \lambda_n)\frac{1}{k_n}\left[\frac{\omega(\lambda)}{\lambda - \lambda_n} - (\phi_\lambda)'_\beta\right].$$

Substituting this formula in (2.20) and letting $\lambda \to \lambda_n$, we get

$$\int_a^{h_1} (\phi_{\lambda_n}(x))^2 dx + \int_{h_1}^{h_2} (\phi_{\lambda_n}(x))^2 dx + \int_{h_2}^{h_3} (\phi_{\lambda_n}(x))^2 dx + \int_{h_3}^{b} (\phi_{\lambda_n}(x))^2 dx = \frac{1}{k_n}\left(\omega'(\lambda_n) - (\phi_{\lambda_n})'_\beta\right).$$

Now putting

$$(\phi_{\lambda_n})'_\beta = \frac{1}{k_n}(\chi_{\lambda_n})'_\beta = \frac{\rho}{k_n}$$

in (2.21) seems that $\omega'(\lambda_n) \neq 0$. ■

## 3 Asymptotic approximate formulas of $\omega(\lambda)$ for four distinct cases

We start by proving some lemmas.

**Lemma 3.1.** *Let $\phi(x,\lambda)$ be the solutions of Eq. (1.1) defined in Section 2, and let $\lambda = s^2$. Then the following integral equations hold for $k = 0, 1$ :*

$$\phi_{1\lambda}^{(k)}(x) = \alpha_2 (\cos s(x-a))^{(k)} - \alpha_1 \frac{1}{s} (\sin s(x-a))^{(k)} + \frac{1}{s} \int_a^x (\sin s(x-y))^k q(y) \phi_{1\lambda}(y)\, dy, \tag{3.1$_k$}$$

$$\phi_{2\lambda}^{(k)}(x) = \frac{\gamma_1}{\delta_1} \phi_{1\lambda}(h_1 - 0) (\cos s(x-h_1))^{(k)} + \frac{1}{s} \frac{\gamma_1'}{\delta_1'} \phi_{1\lambda}'(h_1 - 0) (\sin s(x-h_1))^{(k)} + \frac{1}{s} \int_{h_1}^x (\sin s(x-y))^k q(y) \phi_{2\lambda}(y)\, dy, \tag{3.2$_k$}$$

$$\phi_{3\lambda}^{(k)}(x) = \frac{\gamma_2}{\delta_2} \phi_{2\lambda}(h_2 - 0) (\cos s(x-h_2))^{(k)} + \frac{1}{s} \frac{\gamma_2'}{\delta_2'} \phi_{2\lambda}'(h_2 - 0) (\sin s(x-h_2))^{(k)} + \frac{1}{s} \int_{h_2}^x (\sin s(x-y))^k q(y) \phi_{3\lambda}(y)\, dy, \tag{3.3$_k$}$$

$$\phi_{4\lambda}^{(k)}(x) = \frac{\gamma_3}{\delta_3} \phi_{3\lambda}(h_3 - 0) (\cos s(x-h_3))^{(k)} + \frac{1}{s} \frac{\gamma_3'}{\delta_3'} \phi_{3\lambda}'(h_3 - 0) (\sin s(x-h_3))^{(k)} + \frac{1}{s} \int_{h_3}^x (\sin s(x-y))^k q(y) \phi_{4\lambda}(y)\, dy, \tag{3.4$_k$}$$

*where $(\cdot)^{(k)} = \frac{d^k}{dx^k}(\cdot)$.*

**Proof.** It is enough to substitute $s^2\phi_{1\lambda}(y) + \phi_{1\lambda}''(y)$, $s^2\phi_{2\lambda}(y) + \phi_{2\lambda}''(y)$, $s^2\phi_{3\lambda}(y) + \phi_{3\lambda}''(y)$ and $s^2\phi_{4\lambda}(y) + \phi_{4\lambda}''(y)$ instead of $q(y)\phi_{1\lambda}(y)$, $q(y)\phi_{2\lambda}(y)$, $q(y)\phi_{3\lambda}(y)$, and $q(y)\phi_{4\lambda}(y)$ in the integral terms of the $(3.1_k)$, $(3.2_k)$, $(3.3_k)$ and $(3.4_k)$, respectively, and integrate by parts twice. ■

**Lemma 3.2.** *Let $\lambda = s^2$,* $\operatorname{Im} s = t$. *Then the functions $\phi_{i\lambda}(x)$ have the following asymptotic formulas for $|\lambda| \to \infty$, which hold uniformly for $x \in \Omega_i$*

$(for\ \ i=1,2,3,4.):$

$$\phi_{1\lambda}^{(k)}(x)=\alpha_2\left(\cos s\left(x-a\right)\right)^{(k)}+O\left(\left|s\right|^{k-1}e^{\left|t\right|(x-a)}\right) \qquad (3.5_k)$$

$$\begin{aligned}\phi_{2\lambda}^{(k)}(x)&=\alpha_2\left[\frac{\gamma_1}{\delta_1}\left(\cos s\left(x-h_1\right)\right)^{(k)}\cos s\left(h_1-a\right)-\frac{\gamma_1^{'}}{\delta_1^{'}}\left(\sin s\left(x-h_1\right)\right)^{(k)}\sin s\left(h_1-a\right)\right]\\&\quad+O\left(\left|s\right|^{k-1}e^{\left|t\right|\left[(x-h_1)+(h_1-a)\right]}\right) \qquad (3.6_k)\end{aligned}$$

$$\begin{aligned}\phi_{3\lambda}^{(k)}(x)&=\alpha_2\left\{\frac{\gamma_2}{\delta_2}\left(\cos s\left(x-h_2\right)\right)^{(k)}\left[\frac{\gamma_1}{\delta_1}\cos s\left(h_2-h_1\right)\cos s\left(h_1-a\right)\right.\right.\\&\quad\left.-\frac{\gamma_1^{'}}{\delta_1^{'}}\sin s\left(h_2-h_1\right)\sin s\left(h_1-a\right)\right]-\frac{\gamma_2^{'}}{\delta_2^{'}}\left(\sin s\left(x-h_2\right)\right)^{(k)}\left[\frac{\gamma_1}{\delta_1}\sin s\left(h_2-h_1\right)\cos s\left(h_1-a\right)\right.\\&\qquad (3.7_k)\\&\quad\left.\left.+\frac{\gamma_1^{'}}{\delta_1^{'}}\cos s\left(h_2-h_1\right)\sin s\left(h_1-a\right)\right]\right\}+O\left(\left|s\right|^{k-1}e^{\left|t\right|\left[(x-h_2)+(h_2-h_1)+(h_1-a)\right]}\right)\end{aligned}$$

$$\begin{aligned}\phi_{4\lambda}^{(k)}(x)&=\alpha_2\left\{\frac{\gamma_3}{\delta_3}\left(\cos s\left(x-h_3\right)\right)^{(k)}\left\{\frac{\gamma_2}{\delta_2}\cos s\left(h_3-h_2\right)\left[\frac{\gamma_1}{\delta_1}\cos s\left(h_2-h_1\right)\cos s\left(h_1-a\right)\right.\right.\right.\\&\quad\left.-\frac{\gamma_1^{'}}{\delta_1^{'}}\sin s\left(h_2-h_1\right)\sin s\left(h_1-a\right)\right]-\frac{\gamma_2^{'}}{\delta_2^{'}}\left(\sin s\left(h_3-h_2\right)\right)\left[\frac{\gamma_1}{\delta_1}\sin s\left(h_2-h_1\right)\cos s\left(h_1-a\right)\right.\\&\quad\left.\left.+\frac{\gamma_1^{'}}{\delta_1^{'}}\cos s\left(h_2-h_1\right)\sin s\left(h_1-a\right)\right]\right\}-\frac{\gamma_3^{'}}{\delta_3^{'}}\left(\sin s\left(x-h_3\right)\right)^{(k)}\left\{\frac{\gamma_2}{\delta_2}\sin s\left(h_3-h_2\right)\right.\\&\quad\times\left[\frac{\gamma_1}{\delta_1}\cos s\left(h_2-h_1\right)\cos s\left(h_1-a\right)-\frac{\gamma_1^{'}}{\delta_1^{'}}\sin s\left(h_2-h_1\right)\sin s\left(h_1-a\right)\right]+\frac{\gamma_2^{'}}{\delta_2^{'}}\\&\quad\left.\left.\times\left(\cos s\left(h_3-h_2\right)\right)\left[\frac{\gamma_1}{\delta_1}\sin s\left(h_2-h_1\right)\cos s\left(h_1-a\right)+\frac{\gamma_1^{'}}{\delta_1^{'}}\cos s\left(h_2-h_1\right)\sin s\left(h_1-a\right)\right]\right\}\right\}\\&\quad+O\left(\left|s\right|^{k-1}e^{\left|t\right|\left[(x-h_3)+(h_3-h_2)+(h_2-h_1)+(h_1-a)\right]}\right) \qquad (3.8_k)\end{aligned}$$

*if* $\alpha_2 \neq 0$,

$$\phi_{1\lambda}^{(k)}(x) = -\frac{\alpha_1}{s}\left(\sin s(x-a)\right)^{(k)} + O\left(|s|^{k-2} e^{|t|(x-a)}\right) \tag{$3.9_k$}$$

$$\begin{aligned}\phi_{2\lambda}^{(k)}(x) = & -\frac{\alpha_1}{s}\left[\frac{\gamma_1}{\delta_1}\left(\cos s(x-h_1)\right)^{(k)} \sin s(h_1-a) - \frac{\gamma_1^{'}}{\delta_1^{'}}\left(\sin s(x-h_1)\right)^{(k)} \cos s(h_1-a)\right] \\ & + O\left(|s|^{k-2} e^{|t|[(x-h_1)+(h_1-a)]}\right)\end{aligned} \tag{$3.10_k$}$$

$$\begin{aligned}\phi_{3\lambda}^{(k)}(x) = & -\frac{\alpha_1}{s}\left\{\frac{\gamma_2}{\delta_2}\left(\cos s(x-h_2)\right)^{(k)}\left[\frac{\gamma_1}{\delta_1}\cos s(h_2-h_1)\sin s(h_1-a)\right.\right. \\ & \left.-\frac{\gamma_1^{'}}{\delta_1^{'}}\sin s(h_2-h_1)\cos s(h_1-a)\right] - \frac{\gamma_2^{'}}{\delta_2^{'}}\left(\sin s(x-h_2)\right)^{(k)}\left[\frac{\gamma_1}{\delta_1}\sin s(h_2-h_1)\sin s(h_1-a)\right. \\ & \left.\left.+\frac{\gamma_1^{'}}{\delta_1^{'}}\cos s(h_2-h_1)\cos s(h_1-a)\right]\right\} + O\left(|s|^{k-2} e^{|t|[(x-h_2)+(h_2-h_1)+(h_1-a)]}\right)\end{aligned} \tag{$3.11_k$}$$

$$\begin{aligned}\phi_{4\lambda}^{(k)}(x) = & -\frac{\alpha_1}{s}\left\{\frac{\gamma_3}{\delta_3}\left(\cos s(x-h_3)\right)^{(k)}\left\{\frac{\gamma_2}{\delta_2}\cos s(h_3-h_2)\left[\frac{\gamma_1}{\delta_1}\cos s(h_2-h_1)\sin s(h_1-a)\right.\right.\right. \\ & \left.+\frac{\gamma_1^{'}}{\delta_1^{'}}\sin s(h_2-h_1)\cos s(h_1-a)\right] - \frac{\gamma_2^{'}}{\delta_2^{'}}\sin s(h_3-h_2)\left[\frac{\gamma_1}{\delta_1}\sin s(h_2-h_1)\sin s(h_1-a)\right. \\ & \left.\left.-\frac{\gamma_1^{'}}{\delta_1^{'}}\cos s(h_2-h_1)\cos s(h_1-a)\right]\right\} - \frac{\gamma_3^{'}}{\delta_3^{'}}\left(\sin s(x-h_3)\right)^{(k)}\left\{\frac{\gamma_2}{\delta_2}\sin s(h_3-h_2)\right. \\ & \times\left[\frac{\gamma_1}{\delta_1}\cos s(h_2-h_1)\sin s(h_1-a) + \frac{\gamma_1^{'}}{\delta_1^{'}}\sin s(h_2-h_1)\cos s(h_1-a)\right] - \frac{\gamma_2^{'}}{\delta_2^{'}} \\ & \left.\left.\times\cos s(h_3-h_2)\left[\frac{\gamma_1}{\delta_1}\sin s(h_2-h_1)\sin s(h_1-a) - \frac{\gamma_1^{'}}{\delta_1^{'}}\cos s(h_2-h_1)\cos s(h_1-a)\right]\right\}\right\} \\ & + O\left(|s|^{k-2} e^{|t|[(x-h_3)+(h_3-h_2)+(h_2-h_1)+(h_1-a)]}\right)\end{aligned} \tag{$3.12_k$}$$

*if* $\alpha_2 = 0$.

**Proof.** Since the proof of the formulae for $\phi_{1\lambda}(x)$ is identical to Titchmarsh's proof to similar results for $\phi_\lambda(x)$ (see [24], Lemma 1.7 p. 9-10), we may formulate them without proving them here.

Since the proof of the formulae for $\phi_{2\lambda}(x)$ and $\phi_{3\lambda}(x)$ are identical to Kadakal's and Mukhtarov's proof to similar results for $\phi_\lambda(x)$ (see [3], Lemma 3.2 p. 1373-1375), we may formulate them without proving them here. But the similar formulae for $\phi_{4\lambda}(x)$ need individual consideration, since the last solutions are defined by the initial conditions of these special nonstandart forms. We shall only prove the formula $(3.8_k)$ for $k = 0$.

Let $\alpha_2 \neq 0$. Then according to $(3.7_k)$

$$\phi_{3\lambda}(h_3-0) = \alpha_2 \left\{ \frac{\gamma_2}{\delta_2} \cos s(h_3-h_2) \left[ \frac{\gamma_1}{\delta_1} \cos s(h_2-h_1) \cos s(h_1-a) \right.\right.$$
$$\left. -\frac{\gamma_1'}{\delta_1'} \sin s(h_2-h_1) \sin s(h_1-a) \right] - \frac{\gamma_2'}{\delta_2'} \sin s(h_3-h_2) \left[ \frac{\gamma_1}{\delta_1} \sin s(h_2-h_1) \cos s(h_1-a) \right.$$
$$\left.\left. +\frac{\gamma_1'}{\delta_1'} \cos s(h_2-h_1) \sin s(h_1-a) \right] \right\} + O\left( |s|^{-1} e^{|t|[(h_3-h_2)+(h_2-h_1)+(h_1-a)]} \right)$$

and

$$\phi_{3\lambda}'(h_3-0) = \alpha_2 \left\{ -\frac{s\gamma_2}{\delta_2} \sin s(h_3-h_2) \left[ \frac{\gamma_1}{\delta_1} \cos s(h_2-h_1) \cos s(h_1-a) \right.\right.$$
$$\left. -\frac{\gamma_1'}{\delta_1'} \sin s(h_2-h_1) \sin s(h_1-a) \right] - \frac{s\gamma_2'}{\delta_2'} \cos s(h_3-h_2) \left[ \frac{\gamma_1}{\delta_1} \sin s(h_2-h_1) \cos s(h_1-a) \right.$$
$$\left.\left. +\frac{\gamma_1'}{\delta_1'} \cos s(h_2-h_1) \sin s(h_1-a) \right] \right\} + O\left( e^{|t|[(h_3-h_2)+(h_2-h_1)+(h_1-a)]} \right)$$

Substituting these asymptotic expressions into $(3.4_0)$, we get

$$\phi_{4\lambda}(x) = \alpha_2 \left\{ \frac{\gamma_3}{\delta_3} (\cos s(x-h_3)) \left\{ \frac{\gamma_2}{\delta_2} \cos s(h_3-h_2) \left[ \frac{\gamma_1}{\delta_1} \cos s(h_2-h_1) \cos s(h_1-a) \right.\right.\right.$$
$$\left. -\frac{\gamma_1'}{\delta_1'} \sin s(h_2-h_1) \sin s(h_1-a) \right] - \frac{\gamma_2'}{\delta_2'} \sin s(h_3-h_2) \left[ \frac{\gamma_1}{\delta_1} \sin s(h_2-h_1) \cos s(h_1-a) \right.$$
$$\left.\left. +\frac{\gamma_1'}{\delta_1'} \cos s(h_2-h_1) \sin s(h_1-a) \right] \right\} - \frac{\gamma_3'}{\delta_3'} \sin s(x-h_3) \left\{ \frac{\gamma_2}{\delta_2} \sin s(h_3-h_2) \right.$$
$$\times \left[ \frac{\gamma_1}{\delta_1} \cos s(h_2-h_1) \cos s(h_1-a) - \frac{\gamma_1'}{\delta_1'} \sin s(h_2-h_1) \sin s(h_1-a) \right] + \frac{\gamma_2'}{\delta_2'}$$
$$\left.\left. \times \cos s(h_3-h_2) \left[ \frac{\gamma_1}{\delta_1} \sin s(h_2-h_1) \cos s(h_1-a) + \frac{\gamma_1'}{\delta_1'} \cos s(h_2-h_1) \sin s(h_1-a) \right] \right\} \right\}$$
$$+ \frac{1}{s} \int_{h_3}^{x} \sin s(x-y)\, q(y)\, \phi_{4\lambda}(y)\, dy + O\left( |s|^{-1} e^{|t|[(x-h_3)+(h_3-h_2)+(h_2-h_1)+(h_1-a)]} \right) \tag{3.13}$$

Multiplying through by $e^{-|t|[(x-h_3)+(h_3-h_2)+(h_2-h_1)+(h_1-a)]}$, and denoting

$$F_{4\lambda}(x) := e^{-|t|[(x-h_3)+(h_3-h_2)+(h_2-h_1)+(h_1-a)]} \phi_{4\lambda}(x)$$

we have

$$
\begin{aligned}
F_{4\lambda}(x) := {} & \alpha_2 e^{-|t|[(x-h_3)+(h_3-h_2)+(h_2-h_1)+(h_1-a)]} \Bigg\{ \frac{\gamma_3}{\delta_3} \cos s\,(x-h_3) \Bigg\{ \frac{\gamma_2}{\delta_2} \cos s\,(h_3-h_2) \\
& \times \Bigg[ \frac{\gamma_1}{\delta_1} \cos s\,(h_2-h_1) \cos s\,(h_1-a) - \frac{\gamma_1'}{\delta_1'} \sin s\,(h_2-h_1) \sin s\,(h_1-a) \Bigg] - \frac{\gamma_2'}{\delta_2'} \sin s\,(h_3-h_2) \\
& \times \Bigg[ \frac{\gamma_1}{\delta_1} \sin s\,(h_2-h_1) \cos s\,(h_1-a) + \frac{\gamma_1'}{\delta_1'} \cos s\,(h_2-h_1) \sin s\,(h_1-a) \Bigg] \Bigg] \Bigg\} - \frac{\gamma_3'}{\delta_3'} \sin s\,(x-h_3) \\
& \times \Bigg\{ \frac{\gamma_2}{\delta_2} \sin s\,(h_3-h_2) \Bigg[ \frac{\gamma_1}{\delta_1} \cos s\,(h_2-h_1) \cos s\,(h_1-a) - \frac{\gamma_1'}{\delta_1'} \sin s\,(h_2-h_1) \sin s\,(h_1-a) \Bigg] \\
& + \frac{\gamma_2'}{\delta_2'} \cos s\,(h_3-h_2) \Bigg[ \frac{\gamma_1}{\delta_1} \sin s\,(h_2-h_1) \cos s\,(h_1-a) + \frac{\gamma_1'}{\delta_1'} \cos s\,(h_2-h_1) \sin s\,(h_1-a) \Bigg] \\
& + \frac{1}{s} \int\limits_{h_3}^{x} \sin s\,(x-y)\, q(y)\, e^{-|t|[(x-h_3)+(h_3-h_2)+(h_2-h_1)+(h_1-a)]} F_{4\lambda}(y)\, dy + O\left(|s|^{-1}\right).
\end{aligned}
$$

Denoting $M := \max_{x\in[h_3,b]} |F_{4\lambda}(x)|$ from the last formula, it follows that

$$
\begin{aligned}
M(\lambda) \le {} & |\alpha_2| \left( \left|\frac{\gamma_3}{\delta_3}\right| + \left|\frac{\gamma_3'}{\delta_3'}\right| \right) \left( \left|\frac{\gamma_2}{\delta_2}\right| + \left|\frac{\gamma_2'}{\delta_2'}\right| \right) \left( \left|\frac{\gamma_1}{\delta_1}\right| + \left|\frac{\gamma_1'}{\delta_1'}\right| \right) \\
& + \frac{M(\lambda)}{|s|} \int\limits_{h_3}^{b} q(y)\, dy + \frac{M_0}{|s|}
\end{aligned}
$$

for some $M_0 > 0$. From this, it follows that $M(\lambda) = O(1)$ as $\lambda \to \infty$, so

$$
\phi_{4\lambda}(x) = O\left(e^{|t|[(x-h_3)+(h_3-h_2)+(h_2-h_1)+(h_1-a)]}\right).
$$

Substituting this back into the integral on the right side of (3.13) yields ($3.8_0$). The other cases may be considered analogically. ■

**Theorem 3.1.** *Let* $\lambda = s^2$, $t = \operatorname{Im} s$. *Then the characteristic function* $\omega(\lambda)$ *has the following asymptotic formulas*:

*Case 1*: *If* $\beta_2^{'} \neq 0$, $\alpha_2 \neq 0$, *then*

$$\begin{aligned}
\omega_4(\lambda) = {} & s^3 \beta_2^{'} \alpha_2 \left\{ \frac{\gamma_3}{\delta_3} \left( \sin s \left( b - h_3 \right) \right) \left\{ \frac{\gamma_2}{\delta_2} \cos s \left( h_3 - h_2 \right) \left[ \frac{\gamma_1}{\delta_1} \cos s \left( h_2 - h_1 \right) \cos s \left( h_1 - a \right) \right. \right. \right. \\
& \left. - \frac{\gamma_1^{'}}{\delta_1^{'}} \sin s \left( h_2 - h_1 \right) \sin s \left( h_1 - a \right) \right] - \frac{\gamma_2^{'}}{\delta_2^{'}} \sin s \left( h_3 - h_2 \right) \left[ \frac{\gamma_1}{\delta_1} \sin s \left( h_2 - h_1 \right) \cos s \left( h_1 - a \right) \right. \\
& \left. \left. + \frac{\gamma_1^{'}}{\delta_1^{'}} \cos s \left( h_2 - h_1 \right) \sin s \left( h_1 - a \right) \right] \right\} + \frac{\gamma_3^{'}}{\delta_3^{'}} \cos s \left( b - h_3 \right) \left\{ \frac{\gamma_2}{\delta_2} \sin s \left( h_3 - h_2 \right) \right. \\
& \times \left[ \frac{\gamma_1}{\delta_1} \cos s \left( h_2 - h_1 \right) \cos s \left( h_1 - a \right) - \frac{\gamma_1^{'}}{\delta_1^{'}} \sin s \left( h_2 - h_1 \right) \sin s \left( h_1 - a \right) \right] + \frac{\gamma_2^{'}}{\delta_2^{'}} \\
& \left. \left. \times \cos s \left( h_3 - h_2 \right) \left[ \frac{\gamma_1}{\delta_1} \sin s \left( h_2 - h_1 \right) \cos s \left( h_1 - a \right) + \frac{\gamma_1^{'}}{\delta_1^{'}} \cos s \left( h_2 - h_1 \right) \sin s \left( h_1 - a \right) \right] \right\} \right\} \\
& + O\left( |s|^2 e^{|t|[(b-h_3)+(h_3-h_2)+(h_2-h_1)+(h_1-a)]} \right). \qquad (3.14)
\end{aligned}$$

*Case 2*: *If* $\beta_2^{'} \neq 0$, $\alpha_2 = 0$, *then*

$$\begin{aligned}
\omega_4(\lambda) = {} & -s^2 \alpha_1 \beta_2^{'} \left\{ \frac{\gamma_3}{\delta_3} \left( \sin s \left( b - h_3 \right) \right) \left\{ \frac{\gamma_2}{\delta_2} \cos s \left( h_3 - h_2 \right) \left[ \frac{\gamma_1}{\delta_1} \cos s \left( h_2 - h_1 \right) \sin s \left( h_1 - a \right) \right. \right. \right. \\
& \left. + \frac{\gamma_1^{'}}{\delta_1^{'}} \sin s \left( h_2 - h_1 \right) \cos s \left( h_1 - a \right) \right] - \frac{\gamma_2^{'}}{\delta_2^{'}} \sin s \left( h_3 - h_2 \right) \left[ \frac{\gamma_1}{\delta_1} \sin s \left( h_2 - h_1 \right) \sin s \left( h_1 - a \right) \right. \\
& \left. \left. - \frac{\gamma_1^{'}}{\delta_1^{'}} \cos s \left( h_2 - h_1 \right) \cos s \left( h_1 - a \right) \right] \right\} - \frac{\gamma_3^{'}}{\delta_3^{'}} \cos s \left( b - h_3 \right) \left\{ \frac{\gamma_2}{\delta_2} \sin s \left( h_3 - h_2 \right) \right. \\
& \times \left[ \frac{\gamma_1}{\delta_1} \cos s \left( h_2 - h_1 \right) \sin s \left( h_1 - a \right) + \frac{\gamma_1^{'}}{\delta_1^{'}} \sin s \left( h_2 - h_1 \right) \cos s \left( h_1 - a \right) \right] - \frac{\gamma_2^{'}}{\delta_2^{'}} \\
& \left. \left. \times \cos s \left( h_3 - h_2 \right) \left[ \frac{\gamma_1}{\delta_1} \sin s \left( h_2 - h_1 \right) \sin s \left( h_1 - a \right) - \frac{\gamma_1^{'}}{\delta_1^{'}} \cos s \left( h_2 - h_1 \right) \cos s \left( h_1 - a \right) \right] \right\} \right\} \\
& + O\left( |s| \, e^{|t|[(b-h_3)+(h_3-h_2)+(h_2-h_1)+(h_1-a)]} \right). \qquad (3.15)
\end{aligned}$$

*Case 3*: *If* $\beta_2^{'} = 0$, $\alpha_2 \neq 0$, *then*

$$\begin{aligned}
\omega_4(\lambda) = {} & s^2\alpha_2\beta_1^{'}\Bigg\{\frac{\gamma_3}{\delta_3}\left(\cos s\left(b-h_3\right)\right)\Bigg\{\frac{\gamma_2}{\delta_2}\cos s\left(h_3-h_2\right)\left[\frac{\gamma_1}{\delta_1}\cos s\left(h_2-h_1\right)\cos s\left(h_1-a\right)\right.\\
& \left.-\frac{\gamma_1^{'}}{\delta_1^{'}}\sin s\left(h_2-h_1\right)\sin s\left(h_1-a\right)\right]-\frac{\gamma_2^{'}}{\delta_2^{'}}\sin s\left(h_3-h_2\right)\left[\frac{\gamma_1}{\delta_1}\sin s\left(h_2-h_1\right)\cos s\left(h_1-a\right)\right.\\
& \left.\left.+\frac{\gamma_1^{'}}{\delta_1^{'}}\cos s\left(h_2-h_1\right)\cos s\left(h_1-a\right)\right]\right\}-\frac{\gamma_3^{'}}{\delta_3^{'}}\sin s\left(b-h_3\right)\Bigg\{\frac{\gamma_2}{\delta_2}\sin s\left(h_3-h_2\right)\\
& \times\left[\frac{\gamma_1}{\delta_1}\cos s\left(h_2-h_1\right)\cos s\left(h_1-a\right)-\frac{\gamma_1^{'}}{\delta_1^{'}}\sin s\left(h_2-h_1\right)\sin s\left(h_1-a\right)\right]+\frac{\gamma_2^{'}}{\delta_2^{'}}\\
& \times\cos s\left(h_3-h_2\right)\left[\frac{\gamma_1}{\delta_1}\sin s\left(h_2-h_1\right)\cos s\left(h_1-a\right)+\frac{\gamma_1^{'}}{\delta_1^{'}}\cos s\left(h_2-h_1\right)\sin s\left(h_1-a\right)\right]\Bigg\}\Bigg\}\\
& +O\left(|s|\,e^{|t|[(b-h_3)+(h_3-h_2)+(h_2-h_1)+(h_1-a)]}\right). \qquad (3.16)
\end{aligned}$$

*Case 4*: *If* $\beta_2^{'} = 0$, $\alpha_2 = 0$, *then*

$$\begin{aligned}
\omega_4(\lambda) = {} & -s\alpha_1\beta_1^{'}\Bigg\{\frac{\gamma_3}{\delta_3}\left(\cos s\left(b-h_3\right)\right)\Bigg\{\frac{\gamma_2}{\delta_2}\cos s\left(h_3-h_2\right)\left[\frac{\gamma_1}{\delta_1}\cos s\left(h_2-h_1\right)\sin s\left(h_1-a\right)\right.\\
& \left.+\frac{\gamma_1^{'}}{\delta_1^{'}}\sin s\left(h_2-h_1\right)\cos s\left(h_1-a\right)\right]-\frac{\gamma_2^{'}}{\delta_2^{'}}\sin s\left(h_3-h_2\right)\left[\frac{\gamma_1}{\delta_1}\sin s\left(h_2-h_1\right)\sin s\left(h_1-a\right)\right.\\
& \left.\left.-\frac{\gamma_1^{'}}{\delta_1^{'}}\cos s\left(h_2-h_1\right)\cos s\left(h_1-a\right)\right]\right\}-\frac{\gamma_3^{'}}{\delta_3^{'}}\sin s\left(b-h_3\right)\Bigg\{\frac{\gamma_2}{\delta_2}\sin s\left(h_3-h_2\right)\\
& \times\left[\frac{\gamma_1}{\delta_1}\cos s\left(h_2-h_1\right)\sin s\left(h_1-a\right)+\frac{\gamma_1^{'}}{\delta_1^{'}}\sin s\left(h_2-h_1\right)\cos s\left(h_1-a\right)\right]-\frac{\gamma_2^{'}}{\delta_2^{'}}\\
& \times\cos s\left(h_3-h_2\right)\left[\frac{\gamma_1}{\delta_1}\sin s\left(h_2-h_1\right)\sin s\left(h_1-a\right)-\frac{\gamma_1^{'}}{\delta_1^{'}}\cos s\left(h_2-h_1\right)\cos s\left(h_1-a\right)\right]\Bigg\}\Bigg\}\\
& +O\left(e^{|t|[(b-h_3)+(h_3-h_2)+(h_2-h_1)+(h_1-a)]}\right). \qquad (3.17)
\end{aligned}$$

**Proof.** The proof is immediate by substituting $(3.8_k)$ and $(3.12_k)$ into the representation

$$\begin{aligned}
\omega_4(\lambda) &= \lambda\left(\beta_1^{'}\phi_{4\lambda}(b)-\beta_2^{'}\phi_{4\lambda}^{'}(b)\right)+\left(\beta_1\phi_{4\lambda}(b)-\beta_2\phi_{4\lambda}^{'}(b)\right)\\
&= -\lambda\beta_2^{'}\phi_{4\lambda}^{'}(b)+\lambda\beta_1^{'}\phi_{4\lambda}(b)-\beta_2^{'}\phi_{4\lambda}^{'}(b)+\beta_1\phi_{4\lambda}(b). \qquad (3.18)
\end{aligned}$$

■

**Corollary 3.1.** *The eigenvalues of the problem (1.1)-(1.9) are bounded below.*

**Proof.** Putting $s = it$ $(t > 0)$ in the above formulae, it follows that $\omega_4\left(-t^2\right) \to \infty$ as $t \to \infty$. Therefore, $\omega_4(\lambda) \neq 0$ for $\lambda$ negative and sufficiently large. ■

# 4 Asymptotic formulae for eigenvalues and eigenfunctions

Now we can obtain the asymptotic approximation formulae for the eigenvalues of the considered problem (1.1)-(1.9).

Since the eigenvalues coincide with the zeros of the entire function $\omega_4(\lambda)$, it follows that they have no finite limit. Moreover, we know from Corollaries 2.1 and 2.2 that all eigenvalues are real and bounded below. Hence, we may renumber them as $\lambda_0 \leq \lambda_1 \leq \lambda_2 \leq ...$, listed according to their multiplicity.

In this section, for the sake of simplicity we shall assume that $\gamma_i \delta_i^{'} = \gamma_i^{'} \delta_i$ $(i = 1, 2, 3)$.

**Theorem 4.1.** *The eigenvalues* $\lambda_n = s_n^2$, $n = 0, 1, 2, ...$ *of the problem (1.1)-(1.9) have the following asymptotic formulae for* $n \to \infty$ :

*Case 1*: *If* $\beta_2^{'} \neq 0, \alpha_2 \neq 0$, *then*

$$s_n = \frac{\pi\left(n-1\right)}{b-a} + O\left(\frac{1}{n}\right). \tag{4.1}$$

*Case 2*: *If* $\beta_2^{'} \neq 0, \alpha_2 = 0$, *then*

$$s_n = \frac{\pi\left(n-\frac{1}{2}\right)}{b-a} + O\left(\frac{1}{n}\right). \tag{4.2}$$

*Case 3*: *If* $\beta_2^{'} = 0, \alpha_2 \neq 0$, *then*

$$s_n = \frac{\pi\left(n-\frac{1}{2}\right)}{b-a} + O\left(\frac{1}{n}\right). \tag{4.3}$$

*Case 4*: *If* $\beta_2^{'} = 0, \alpha_2 = 0$, *then*

$$s_n = \frac{\pi n}{b-a} + O\left(\frac{1}{n}\right). \tag{4.4}$$

**Proof.** We shall only consider the first case. The other cases may be considered similarly. Denoting $\overline{\omega}(s) := \omega_4\left(s^2\right) = \omega_4(\lambda)$,

$$\overline{\omega}_1(s) = \frac{\alpha_2 \beta_2^{'} \gamma_3 \gamma_2 \gamma_1 s^3}{\delta_3 \delta_2 \delta_1} \sin\left(s\left(b-a\right)\right)$$

and $\overline{\omega}_2(s) := \overline{\omega}(s) - \overline{\omega}_1(s)$, we write $\overline{\omega}(s)$ as $\overline{\omega}(s) = \overline{\omega}_1(s)$. In view of (3.14), from basic properties of trigonometric functions we have

$$\overline{\omega}_2(s) = O\left(\left|s\right|^2 e^{\left|t\right|\left(b-a\right)}\right).$$

We shall apply the well-known Rouché's theorem, which asserts that if $f(s)$ and $g(s)$ are analytic inside and on a closed contour $C$, and $\left|g(s)\right| < \left|f(s)\right|$ on $C$, then $f(s)$ and $f(s) + g(s)$ have the same number zeros inside $C$, provided that each zero is counted according to their multiplicity. It is readily shown that $|\overline{\omega}_1(s)| > |\overline{\omega}_2(s)|$ on the contours

$$C_n := \left\{ s \in \mathbb{C} \, | |s| = \frac{\left(n + \frac{1}{2}\right)\pi}{(b-a)} \right\}$$

for sufficiently large $n$.

Let $\lambda_0 \leq \lambda_1 \leq \lambda_2 \leq \ldots$ be zeros of $\omega(\lambda)$ and $\lambda_n = s_n^2$. Since inside the contour $C_n$, $\overline{\omega}_1(s)$ has zeros at points $s = 0$ with multiplicity 4 and $s = \frac{k\pi}{2(b-a)}$, $k = \pm 1, \pm 2, \ldots, \pm n$ with multiplicity 1. The number of zeros is $2n+4$, and it follows that

$$s_n = \frac{(n-1)\pi}{b-a} + \delta_n \tag{4.5}$$

where $\delta_n = O(1)$ for sufficiently large $n$. By substituting this in (3.14), we derive that $\delta_n = O\left(\frac{1}{n}\right)$, which completes the proof. ■

The next approximation for the eigenvalues may be obtained by the following procedure. For this, we shall suppose that $q(y)$ is of bounded variation in $[a,b]$.

We only consider the case $\beta_2^{'} \neq 0$ and $\alpha_2 \neq 0$ (since the other cases may be considered anologically). Putting $x = h_1$ in $(3.1_k)$, $x = h_2$ in $(3.2_k)$, $x = h_3$ in $(3.3_k)$ and then substituting in $(3.4_k)$, we get that

$$\begin{aligned}
\phi_{4\lambda}^{'}(b) &= s\frac{\alpha_2\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3}\sin(s(b-a)) - \frac{\alpha_1\gamma_1^{'}\gamma_2^{'}\gamma_3^{'}}{\delta_1^{'}\delta_2^{'}\delta_3^{'}}\cos(s(b-a)) \\
&+ \frac{\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3}\int_a^{h_1}\cos(s(b-y))\,q(y)\,\phi_{1\lambda}(y)\,dy \\
&+ \frac{\gamma_2\gamma_3}{\delta_2\delta_3}\int_{h_1}^{h_2}\cos(s(b-y))\,q(y)\,\phi_{2\lambda}(y)\,dy \\
&+ \frac{\gamma_3}{\delta_3}\int_{h_2}^{h_3}\cos(s(b-y))\,q(y)\,\phi_{3\lambda}(y)\,dy + \int_{h_3}^{b}\cos(s(b-y))\,q(y)\,\phi_{4\lambda}(y)\,dy.
\end{aligned}$$

Substituting $(3.5_k)$, $(3.6_k)$, $(3.7_k)$ and $(3.8_k)$ into the right side of the last integral equality then gives

$$\begin{aligned}\phi_{4\lambda}'(b) &= -s\frac{\alpha_2\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3}\sin(s(b-a)) - \frac{\alpha_1\gamma_1'\gamma_2'\gamma_3'}{\delta_1'\delta_2'\delta_3'}\cos(s(b-a)) \\ &\quad + \frac{\alpha_2\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3}\int_a^{h_1}\cos(s(b-y))\cos(s(y-a))\,q(y)\,dy + \frac{\alpha_2\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3} \\ &\quad \times \int_{h_1}^{h_2}\cos(s(b-y))\cos(s(y-a))\,q(y)\,dy + \frac{\alpha_2\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3} \\ &\quad \times \int_{h_2}^{h_3}\cos(s(b-y))\cos(s(y-a))\,q(y)\,dy + \frac{\alpha_2\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3} \\ &\quad \times \int_{h_3}^{b}\cos(s(b-y))\cos(s(y-a))\,q(y)\,dy + O\left(|s|^{-1}e^{|t|[b-a]}\right).\end{aligned}$$

On the other hand, from $(3.8_k)$, it follows that

$$\phi_{4\lambda}(b) = \frac{\alpha_2\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3}\cos(s(b-a)) + O\left(|s|^{-1}e^{|t|[b-a]}\right)$$

Putting these formulas into (3.18), we have

$$\begin{aligned}\omega_4(\lambda) &= s^3\frac{\beta_2'\alpha_2\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3}\sin(s(b-a)) + s^2\left[\left(\frac{\beta_1'\alpha_2\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3} + \frac{\beta_2'\alpha_1\gamma_1'\gamma_2'\gamma_3'}{\delta_1'\delta_2'\delta_3'}\right)\right. \\ &\quad \times\cos(s(b-a)) - \frac{\beta_2'\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3}\int_a^{h_1}\cos(s(b-y))\,q(y)\,\phi_{1\lambda}(y)\,dy \\ &\quad - \frac{\beta_2'\gamma_2\gamma_3}{\delta_2\delta_3}\int_{h_1}^{h_2}\cos(s(b-y))\,q(y)\,\phi_{2\lambda}(y)\,dy - \frac{\beta_2'\gamma_3}{\delta_3}\int_{h_2}^{h_3}\cos(s(b-y))\,q(y)\,\phi_{3\lambda}(y)\,dy \\ &\quad \left. -\beta_2'\int_{h_3}^{b}\cos(s(b-y))\,q(y)\,\phi_{4\lambda}(y)\,dy\right] + O\left(|s|^{-1}e^{|t|[b-a]}\right).\end{aligned}$$

Putting (4.5) in the last equality we find that

$$\delta_n = -\frac{1}{s_n}\left[\frac{\beta_1'}{\beta_2'} + \frac{\alpha_1}{\alpha_2} - R + O\left(\frac{1}{n}\right)\right] + O\left(|s_n|^{-2}\right) \tag{4.6}$$

where $R := \frac{1}{2}\left[\int_a^{h_1}q(y)\,dy + \int_{h_1}^{h_2}q(y)\,dy + \int_{h_2}^{h_3}q(y)\,dy + \int_{h_3}^{b}q(y)\,dy\right]$. Recalling that $q(y)$ is of bounded variation in $[a,b]$, and applying the well-known Riemann-

Lebesque Lemma (see [25], p. 48, Theorem 4.12) to the fourth integral on the right in (4.6), this term is $O\left(\frac{1}{n}\right)$. As a result, from (4.6) it follows that

$$\delta_n = -\frac{(b-a)}{\pi(n-1)}\left[\frac{\beta_1'}{\beta_2'} + \frac{\alpha_1}{\alpha_2} - R\right] + O\left(\frac{1}{n^2}\right).$$

Substituting in (4.5), we have

$$s_n = \frac{\pi(n-1)}{b-a} - \frac{(b-a)}{\pi(n-1)}\left[\frac{\beta_1'}{\beta_2'} + \frac{\alpha_1}{\alpha_2} - R\right] + O\left(\frac{1}{n^2}\right).$$

Similar results in the other cases are as follows:
In case 2:

$$s_n = \frac{\pi\left(n-\frac{1}{2}\right)}{b-a} + \frac{1}{\pi\left(n-\frac{1}{2}\right)}\left[\frac{\beta_1'}{\beta_2'} - R\right] + O\left(\frac{1}{n^2}\right).$$

In case 3:

$$s_n = \frac{\pi\left(n-\frac{1}{2}\right)}{b-a} + \frac{1}{\pi\left(n-\frac{1}{2}\right)}\left[-\frac{\beta_2}{\beta_1'} + \frac{\alpha_1}{\alpha_2} - R\right] + O\left(\frac{1}{n^2}\right).$$

In case 4:

$$s_n = \frac{\pi n}{b-a} + \frac{1}{\pi n}\left[\frac{\beta_2}{\beta_1'} + R\right] + O\left(\frac{1}{n^2}\right).$$

By putting (4.1) into the ($3.5_k$), ($3.6_k$), ($3.7_k$) and ($3.8_k$) we obtain that

$$\phi_{1\lambda_n}(x) = \alpha_2 \cos\left(\frac{\pi(n-1)(x-a)}{b-a}\right) + O\left(\frac{1}{n}\right),$$

$$\phi_{2\lambda_n}(x) = \frac{\alpha_2\gamma_1}{\delta_1}\cos\left(\frac{\pi(n-1)(x-a)}{b-a}\right) + O\left(\frac{1}{n}\right),$$

$$\phi_{3\lambda_n}(x) = \frac{\alpha_2\gamma_1\gamma_2}{\delta_1\delta_2}\cos\left(\frac{\pi(n-1)(x-a)}{b-a}\right) + O\left(\frac{1}{n}\right)$$

and

$$\phi_{4\lambda_n}(x) = \frac{\alpha_2\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3}\cos\left(\frac{\pi(n-1)(x-a)}{b-a}\right) + O\left(\frac{1}{n}\right)$$

in the first case. Consequently, if $\beta_2' \neq 0$ and $\alpha_2 \neq 0$, then the eigenfunction $\phi(x, \lambda_n)$ has the following asymptotic formulae

$$\phi(x,\lambda_n) = \begin{cases} \alpha_2\cos\left(\frac{\pi(n-1)(x-a)}{b-a}\right) + O\left(\frac{1}{n}\right), & x \in [a, h_1) \\ \frac{\alpha_2\gamma_1}{\delta_1}\cos\left(\frac{\pi(n-1)(x-a)}{b-a}\right) + O\left(\frac{1}{n}\right), & x \in (h_1, h_2) \\ \frac{\alpha_2\gamma_1\gamma_2}{\delta_1\delta_2}\cos\left(\frac{\pi(n-1)(x-a)}{b-a}\right) + O\left(\frac{1}{n}\right), & x \in (h_2, h_3) \\ \frac{\alpha_2\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3}\cos\left(\frac{\pi(n-1)(x-a)}{b-a}\right) + O\left(\frac{1}{n}\right), & x \in (h_3, b] \end{cases}$$

which holds uniformly for $x \in [a, h_1) \cup (h_1, h_2) \cup (h_2, h_3) \cup (h_3, b]$.

Similar formulae in the other cases are as follows:

In case 2

$$\phi\left(x,\lambda_n\right)=\begin{cases} -\frac{\alpha_1(b-a)}{\pi\left(n-\frac{1}{2}\right)}\sin\left(\frac{\pi\left(n-\frac{1}{2}\right)(x-a)}{b-a}\right)+O\left(\frac{1}{n^2}\right), & x\in[a,h_1) \\ -\frac{\alpha_1(b-a)\gamma_1}{\delta_1\pi\left(n-\frac{1}{2}\right)}\sin\left(\frac{\pi\left(n-\frac{1}{2}\right)(x-a)}{b-a}\right)+O\left(\frac{1}{n^2}\right), & x\in(h_1,h_2) \\ -\frac{\alpha_1(b-a)\gamma_1\gamma_2}{\delta_1\delta_2\pi\left(n-\frac{1}{2}\right)}\sin\left(\frac{\pi\left(n-\frac{1}{2}\right)(x-a)}{b-a}\right)+O\left(\frac{1}{n^2}\right), & x\in(h_2,h_3) \\ -\frac{\alpha_1(b-a)\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3\pi\left(n-\frac{1}{2}\right)}\sin\left(\frac{\pi\left(n-\frac{1}{2}\right)(x-a)}{b-a}\right)+O\left(\frac{1}{n^2}\right), & x\in(h_3,b]\,. \end{cases}$$

In case 3

$$\phi\left(x,\lambda_n\right)=\begin{cases} \alpha_2\cos\left(\frac{\pi\left(n-\frac{1}{2}\right)(x-a)}{b-a}\right)+O\left(\frac{1}{n}\right), & x\in[a,h_1) \\ \frac{\alpha_2\gamma_1}{\delta_1}\cos\left(\frac{\pi\left(n-\frac{1}{2}\right)(x-a)}{b-a}\right)+O\left(\frac{1}{n}\right), & x\in(h_1,h_2) \\ \frac{\alpha_2\gamma_1\gamma_2}{\delta_1\delta_2}\cos\left(\frac{\pi\left(n-\frac{1}{2}\right)(x-a)}{b-a}\right)+O\left(\frac{1}{n}\right), & x\in(h_2,h_3) \\ \frac{\alpha_2\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3}\cos\left(\frac{\pi\left(n-\frac{1}{2}\right)(x-a)}{b-a}\right)+O\left(\frac{1}{n}\right), & x\in(h_3,b]\,. \end{cases}$$

In case 4

$$\phi\left(x,\lambda_n\right)=\begin{cases} -\frac{\alpha_1(b-a)}{\pi n}\sin\left(\frac{\pi n(x-a)}{b-a}\right)+O\left(\frac{1}{n^2}\right), & x\in[a,h_1) \\ -\frac{\alpha_1(b-a)\gamma_1}{\delta_1\pi n}\sin\left(\frac{\pi n(x-a)}{b-a}\right)+O\left(\frac{1}{n^2}\right), & x\in(h_1,h_2) \\ -\frac{\alpha_1(b-a)\gamma_1\gamma_2}{\delta_1\delta_2\pi n}\sin\left(\frac{\pi n(x-a)}{b-a}\right)+O\left(\frac{1}{n^2}\right), & x\in(h_2,h_3) \\ -\frac{\alpha_1(b-a)\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3\pi n}\sin\left(\frac{\pi n(x-a)}{b-a}\right)+O\left(\frac{1}{n^2}\right), & x\in(h_3,b]\,. \end{cases}$$

All these asymptotic formulas hold uniformly for $x \in [a, h_1) \cup (h_1, h_2) \cup (h_2, h_3) \cup (h_3, b]$.

# References

[1] M. Demirci , Z. Akdoğan, O.Sh. Mukhtarov, Asymptotic behavior of eigenvalues and eigenfuctions of one discontinuous boundary-value problem, Intenational Journal of Computational Cognition 2(3) (2004) 101–113.

[2] D. Buschmann, G. Stolz, J. Weidmann, One-dimensional Schrödinger operators with local point interactions, Journal für die Reine und Angewandte Mathematik 467 (1995) 169–186.

[3] M. Kadakal, O.Sh. Mukhtarov, Sturm-Liouville problems with discontinuities at two points, Comput. Math. Appl.,54 (2007) 1367-1379.

[4] Q. Yang, W. Wang, Asymptotic behavior of a differential operator with discontinuities at two points, Math. Meth. Appl. Sci. 34 (2011) 373-383.

[5] M. Kadakal, O.Sh. Mukhtarov, F.Ş. Muhtarov, Some spectral problems of Sturm-Liouville problem with transmission conditions, Iranian Journal of Science and Technology, 49(A2) (2005) 229-245.

[6] Z. Akdoğan, M. Demirci, O.Sh. Mukhtarov, Green function of discontinuous boundary-value problem with transmission conditions. Mathematical Models and Methods in Applied Sciences 30 (2007) 1719-1738.

[7] C.T. Fulton, Two-point boundary value problems with eigenvalue parameter contained in the boundary conditions, Proc. Roy. Soc. Edinburgh Sect. A 77 (1977) 293–308.

[8] M. Kobayashi, Eigenvalues of discontinuous Sturm–Liouville problems with symmetric potential, Comput. Math. Appl. 18 (4) (1989) 357–364.

[9] A.V. Likov, Yu.A. Mikhailov, The Theory of Heat and Mass Transfer, Qosenerqoizdat (1963) (in Russian).

[10] E. Tunç, O.Sh. Mukhtarov, Fundamental solutions and eigenvalues of one boundary-value problem with transmission conditions, Appl. Math. Comput. 157 (2004) 347–355.

[11] O.Sh. Mukhtarov, E. Tunç, Eigenvalue problems for Sturm–Liouville equations with transmission conditions, Israel J. Math. 144 (2004) 367–380.

[12] O.Sh. Mukhtarov, M. Kadakal, F.Ş. Muhtarov, Eigenvalues and normalized eigenfunctions of discontinuous Sturm–Liouville problem with transmission conditions, Rep. Math. Phys. 54 (2004) 41–56.

[13] P.A. Binding, P.J. Browne, Oscillation theory for indefinite Sturm–Liouville problems with eigen-parameter-dependent boundary conditions, Proc. Roy. Soc. Edinburgh Sect. A 127 (1997) 1123–1136.

[14] D.B. Hinton, An expansion theorem for an eigenvalue problem with eigenvalue parameter in the boundary conditions, Q. J. Math. 30 (1979) 33–42.

[15] N.B. Kerimov, Kh.K. Mamedov, On a boundary value problem with a spectral parameter in the boundary conditions, Sibirsk. Mat. Zh. 40 (2) (1999) 325–335 (English translation: Siberian Math. J. 40 (2) (1999) 281–290).

[16] A.A. Shkalikov, Boundary value problems for ordinary differential equations with a parameter in boundary condition, Trudy Sem. Imeny I. G. Petrowsgo 9 (1983) 190–229.

[17] J. Walter, Regular eigenvalue problems with eigenvalue parameter in the boundary conditions, Math. Z. 133 (1973) 301–312.

[18] S. Yakubov, Y. Yakubov, Abel basis of root functions of regular boundary value problems, Math. Nachr. 197 (1999) 157–187.

[19] S. Yakubov, Y. Yakubov, Differential-Operator Equations. Ordinary and Partial Differential Equations, Chapman and Hall/CRC, Boca Raton, 2000.

[20] E. Şen, A. Bayramov, Calculation of eigenvalues and eigenfunctions of a discontinuous boundary value problem with retarded argument which contains a spectral parameter in the boundary condition, Mathematical and Computer Modelling, 54 (2011) 3090-3097.

[21] O.Sh. Mukhtarov, N. Kuruoglu, M. Kandemir, Distribution of eigenvalues for the discontinuous boundary value problem with functional manypoint conditions, Israel J. Math. 129 (2002) 143–156.

[22] O. Muhtarov, S. Yakubov, Problems for ordinary differential equations with transmission conditions, Appl. Anal. 81(5) (2002) 1033–1064.

[23] M.N. Naimark, Linear Differential Operators, Ungar, New York, 1967.

[24] E.C. Titchmarsh, Eigenfunctions Expansion Associated With Second Order Differential Equation 1, 2nd ed., Oxford University Press, London, 1962.

[25] A. Zygmund, Trigonometric Series I, 2nd ed., Pergamon, London, 1959